

Keep out of reach of children  
FOR ANIMAL TREATMENT ONLY

# ANDROVAX<sup>®</sup>plus

Increase the number of lambs born to breeding ewes

Vaccine for Sheep

Contains Androstenedione-protein complex

200mL and 500mL

Read entire label before use.

**HANDLING PRECAUTIONS WARNING** - may cause an allergic skin reaction. Avoid skin contact.

**FIRST AID.** If on skin, wash with plenty of soap and water. If skin irritation/rash occurs seek medical advice. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor.

**ACCIDENTAL SELF-INJECTION** - Obtain medical attention - show this leaflet and/or SDS Accidental self injection may lead to an inflammatory response and medical advice should be sought on the management of deep injections, particularly those near a joint or associated with bruising. If possible the application of gentle squeezing pressure with absorbent material (e.g. facial tissues) at the injection site will swab up unabsorbed vaccine. Strong squeezing of the site should be avoided. The damaged area should be thoroughly cleansed and a topical antiseptic applied.

**ADVICE TO DOCTOR** - Treat symptomatically. Some risk of hypersensitivity from injection. Contamination of the needle must be considered.

**DISPOSAL** Dispose of contents and empty vaxipak by crushing or puncturing and burying in an approved landfill.

See Safety Data Sheet for further information. [www.intervet.co.nz](http://www.intervet.co.nz).

## VACCINATION PROGRAMME

	<b>Sensitiser</b>	<b>Booster</b>
All breeding sheep in first year of use	8-10 weeks before mating	4-6 weeks before mating, and not less than 4 weeks after 1 <sup>st</sup> dose.
Subsequent years of use	Not required	Not less than 4 weeks, preferably 4-6 weeks before mating

## DOSAGE

Inject a **2mL dose subcutaneously** in the anterior (front) half of the neck.

## DIRECTIONS FOR USE

**Shake well before use.**

### Equipment:

1. Ensure vaccinator delivers a **2mL dose**.
2. Attach end of draw-off tube to vaccinator. Pierce rubber bung of Vaxipak with draw-off needle.
3. Prime vaccinator as per vaccinator instructions.
4. Use sharp needles. Change needles approx every 20 sheep.
5. Attach tie to vaxipack, and hang around neck.
6. Needles and vaccinator should be sterile prior to use.

### Vaccination Technique:

1. **Vaccinate only healthy, clean, dry sheep.**
2. Inject in the anterior (front) half of the neck.
3. Inject **subcutaneously** i.e. **between the skin and muscle**.
4. Part wool, raise skin to form a "tent" and insert needle into "tent opening" so that the needle is almost parallel with the neck.

## PRECAUTIONS

- A slight reaction may occur at the site of injection but will normally disappear in a few weeks
- Ewes may not show signs of oestrus and may be temporarily infertile for 2-3 weeks following the second or booster vaccination (see Vaccination Programme).

## STORAGE

Store away from light at 2-8°C. **DO NOT FREEZE.**

Unused vaccine must be discarded within 12 hours of opening.

Prescription Animal Remedy (**P.A.R**) Class I.

For use only under the authority or prescription of a veterinarian.

Registered pursuant to the ACVM Act 1997, No. A9927

See [www.nzfsa.govt.nz/acvm/](http://www.nzfsa.govt.nz/acvm/) for registration conditions

Registered pursuant to the HSNO Act 1996, Approval code HSR000015

See [www.ermanz.govt.nz](http://www.ermanz.govt.nz) for controls

ANDROVAX and VAXIPAK are registered trademarks of Schering-Plough Animal Health Ltd.

Registered to:

**Schering-Plough Animal Health Ltd**

Upper Hutt, New Zealand

Phone: 0800 800 543

[www.intervet.co.nz](http://www.intervet.co.nz)