

RESTRICTED VETERINARY MEDICINE

Keep out of reach of Children

FOR ANIMAL TREATMENT ONLY

Bovaclox™

dry cow

20 x Syringes

Each single dose syringe contains 500mg cloxacillin (as the benzathine salt) and 250mg ampicillin (as the trihydrate) in a long acting base.

Read entire label before use

HANDLING PRECAUTIONS WARNING Contains penicillins which may cause skin irritation or an allergic skin reaction. Do not handle this product if you know you are sensitised, or if you have been advised not to work with such preparations. Use gloves when handling this product to avoid exposure, wash hands after use.

FIRST AID In case of skin contact, carefully remove any contaminated clothing and wash skin thoroughly with soap and water. If irritation or symptoms occur or persist, consult a doctor. Swelling of the face, eyes or lips or difficulty with breathing are more serious symptoms and require urgent medical attention. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor.

DISPOSAL Dispose of unused contents in a suitable landfill. Dispose of empty syringes in a suitable landfill. Do NOT burn.

See Safety Data Sheet for further information. www.msd-animal-health.co.nz

USES

Routine dry cow therapy is an integral part of mastitis control. In conjunction with teat spraying and proper management of the cow during the drying off period, the careful administration of dry cow therapy reduces new infections at drying off and in the early dry period. Dry cow therapy is also useful in treating subclinical mastitis that may be present at drying off.

PRECAUTIONS

Not for use in lactating dairy cows. Dry cow therapy should be used once at drying off only. If mastitis develops during the dry period a lactating cow product should be used.

STORAGE

Store protected from light between 5-30 °C. Do not refrigerate or freeze.

DOSAGE AND ADMINISTRATION

Treatment is to be at least 30 days before calving.

When using dry cow preparations, care must be taken not to introduce infection into the udder.

1. After milking is complete, thoroughly clean and disinfect the end of the teat (e.g. with a fresh teatwipe).
2. Hold the barrel of the syringe firmly in one hand and gently twist and pull the protective cap to remove it. DO NOT BEND THE NOZZLE. Take care not to contaminate the nozzle.
3. Partially insert the nozzle into the teat canal and apply steady pressure on the syringe plunger until the full dose has been delivered.
4. Finally, spray carefully with an approved teat spray.

WITHHOLDING PERIOD

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residue Limits of Agricultural Compounds) Food Standards.

Milk: Treatment is to be at least **30** days before calving.

Milk (colostrum) from the first 8 milkings after calving should be prevented from directly entering the human food chain. If calving occurs within 30 days of the last treatment, milk sold for human consumption may be taken only after the full 30 days from treatment and a further 8 milkings have elapsed.

Meat: Animals producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 30 days of the last treatment.

ACVM Registration No: A4495.

See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to the HSNO Act 1996, HSR002346.

See www.epa.govt.nz for controls.

Bovaclox is a trademark. ©2012

Registered to:

Schering-Plough Animal Health Limited,

Phone: 0800 800 543

www.msd-animal-health.co.nz

