

Keep out of reach of children
FOR ANIMAL TREATMENT ONLY
CAMPYVAX4[®]

Prevents abortion and perinatal loss caused by *Campylobacter fetus fetus*
Induces a specific immune response to aid the protection against abortion
due to *C. jejuni*

Contains inactivated *Campylobacter fetus* subspecies *fetus*, serotypes
DL42, 6/1, 134 and *Campylobacter jejuni*.

100mL and 500mL

Read entire label before use.

HANDLING PRECAUTION WARNING may cause an allergic skin reaction. Avoid skin contact.

FIRST AID. If on skin, wash with plenty of soap and water. If skin irritation/rash occurs seek medical advice. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor.

ACCIDENTAL SELF-INJECTION - Obtain medical attention - show this leaflet and/or SDS Accidental self injection may lead to an inflammatory response and medical advice should be sought on the management of deep injections, particularly those near a joint or associated with bruising. If possible the application of gentle squeezing pressure with absorbent material (e.g. facial tissues) at the injection site will swab up unabsorbed vaccine. Strong squeezing of the site should be avoided. The damaged area should be thoroughly cleansed and a topical antiseptic applied.

ADVICE TO DOCTOR - Treat symptomatically. Some risk of hypersensitivity from injection. Contamination of the needle must be considered.

DISPOSAL Dispose of contents and empty vaxipak by burying in an approved landfill.

See Safety Data Sheet for further information. www.intervet.co.nz.

VACCINATION PROGRAMME

	Sensitiser	Booster
All breeding sheep in first year of use	Inject pre-tup	Inject 4-8 weeks later (ideally pre-tup)
Subsequent years of use	Not required	Inject pre-tup

DOSAGE

Inject a **1mL dose subcutaneously** in the anterior (front) half of the neck.

DIRECTIONS FOR USE

Shake well.

Equipment:

1. Ensure vaccinator delivers a **1mL dose**.
2. Attach end of draw-off tube to vaccinator. Pierce rubber bung of vaxipak with draw-off needle.

3. Prime vaccinator as per vaccinator instructions.
4. Use sharp needles. Change needles approx every 20 sheep.
5. Needles and vaccinator should be sterile prior to use.

Vaccination Technique:

1. **Vaccinate only healthy, clean, dry sheep.**
2. Inject in the anterior (front) half of the neck.
3. Inject subcutaneously i.e. between the skin and muscle.
4. Part wool, raise skin to form a "tent" and insert needle into "tent opening" so that the needle is almost parallel with the neck.

PRECAUTIONS

- A slight reaction may occur at the site of injection but will normally disappear in a few weeks
- Occasionally following vaccination some ewes may develop symptoms such as lethargy, mild loss of appetite and ataxia.

STORAGE

Store away from light at 2-8°C. Do not freeze. Unused vaccine must be discarded within 12 hours of opening.

Prescription Animal Remedy (**P.A.R**) Class I.

For use only under the authority or prescription of a veterinarian.

Registered pursuant to the ACVM Act 1997, No. A9535

See www.nzfsa.govt.nz/acvm/ for registration conditions

Approved pursuant to the HSNO Act 1996, Approval Code HSR000015.

CAMPYVAX4 and VAXIPAK are registered trademarks of Schering-Plough Animal Health Ltd.

Registered to:

Schering-Plough Animal Health Ltd

33 Whakatiki St,

Upper Hutt, New Zealand

Phone: 0800 800 543