

Keep out of reach of children  
FOR ANIMAL TREATMENT ONLY

# CEFA-SAFE™

Dry Cow Preparation

Active Ingredient: cephapirin 300mg/10mL syringe

**200 SYRINGES (50 COWS)**

## READ ENTIRE LABEL BEFORE USE

**HANDLING PRECAUTIONS** Warning. May cause mild skin irritation and/or an allergic skin reaction. May cause organ damage from repeated oral exposure at high doses. Do not handle this product if you know you are sensitised, or if you have been advised not to work with such preparations. Use gloves when handling this product to avoid exposure, wash hands after use.

**FIRST AID** If on skin, carefully remove any contaminated clothing and wash with plenty of soap and water. If skin irritation or rash occurs, get medical advice.

**DISPOSAL** Dispose of empty containers by burying in a suitable landfill.

See safety data sheet for further information. [www.intervet.co.nz](http://www.intervet.co.nz)

## INDICATIONS

Treatment of subclinical mastitis and prevention of mastitis caused by bacteria sensitive to cephapirin in cows at drying off.

## DESCRIPTION

Intramammary infusion consisting of a 10mL syringe containing 300mg cephapirin as cephapirin benzathine.

## MODE OF ACTION

Cephapirin is an antibiotic of the  $\beta$ -lactam group of cephalosporins, and is active against many Gram-positive and some Gram-negative organisms.

## CONTRAINDICATIONS

Not to be used in animals known to be allergic to cephalosporins.


## DOSAGE AND ADMINISTRATION

### Dosage

The contents of one syringe are infused into each quarter when the cow is dried off. Dry cow therapy should be used at drying off only.

### Administration

Treat each cow immediately following its final milking for the season. Administration must not be delayed. Before use, milk the udder completely out. Care must be taken not to introduce infection into the udder. Clean the teat thoroughly with cotton wool soaked in a 70% alcohol solution and allow to dry. Remove the cap from the end of the nozzle of the syringe partially or totally, depending on partial or total insertion. Insert the nozzle carefully into the opening of the teat, and infuse the full contents of the syringe slowly into the teat canal. Gently massage the quarter to disperse the suspension upwards. Spray carefully with an approved teat spray.



Lactating cow products (e.g. Penalone A7795 or Spectrazol Milking Cow A5270) should be used if retreatment is required during the dry period.

## **WITHHOLDING PERIODS**

**Milk:** Treatment to be at least 35 days before calving. If calving occurs within 35 days of the last treatment, milk to be sold for human consumption may be taken only after the full 35 days from treatment and a further 8 milkings have elapsed. Milk (colostrum) from the first 8 milkings after calving should be prevented from directly entering the human food chain.

**Meat:** Cattle producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 21 days of the last treatment.

## **STORAGE**

Store below 30°C. Do not refrigerate or freeze.

Prescription Animal Remedy (**P.A.R**) Class I.

For use only under the authority or prescription of a veterinarian.

Registered pursuant to the ACVM Act 1997, No. A7158. See [www.nzfsa.govt.nz/acvm/](http://www.nzfsa.govt.nz/acvm/) for registration conditions.

Approved pursuant to the HSNO Act 1996, HSR002350. See [www.ermanz.govt.nz](http://www.ermanz.govt.nz) for controls.

Registered to:

Schering-Plough Animal Health Ltd

33 Whakatiki St, Upper Hutt.

Phone: 0800 800 543