

## RESTRICTED VETERINARY MEDICINE

Keep out of reach of children  
FOR ANIMAL TREATMENT ONLY

# DEPOCILLIN<sup>®</sup>

INJECTABLE AQUEOUS SOLUTION OF PROCAINE PENICILLIN

300,000 IU (300mg) procaine penicillin per mL.

**100mL and 250mL**

### Read the entire label before use

**HANDLING PRECAUTIONS** Warning, contains penicillin which may cause an allergic skin reaction. Do not handle this product if you know you are sensitised, or if you have been advised not to work with such preparations. Use gloves, wash hands after use.

**FIRST AID If on skin** wash thoroughly with soap and water. If irritation or symptoms occur or persist, consult a doctor. Swelling of the face, eyes or lips or difficulty with breathing are more serious symptoms and require urgent medical attention. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately.

**DISPOSAL** Dispose of any unused contents and empty container in an approved landfill. Avoid release to the environment and avoid contamination of any water supply with product or empty container.

See Safety Data Sheet for further information. [www.intervet.co.nz](http://www.intervet.co.nz)

### INDICATIONS

The treatment of infections caused by bacteria sensitive to penicillin in horses, cattle, sheep, pigs, dogs and cats.

### DOSAGE

Large animals 12,000 IU/kg (12mg/kg liveweight) **1mL/25kg by i.m.**

e.g. Horse 500kg 20mL  
Cow 400kg 16mL  
Sheep 50kg 2mL  
Pig 50kg 2mL

Not for use in bobby calves.

Companion animals 30,000 IU/kg (30mg/kg liveweight) **1mL/10kg by s.c.**

e.g. Dog 10kg 1mL  
Cat 5kg 0.5mL

### ADMINISTRATION

In food producing animals the injection is to be given into the anterior half of the neck. Clean the area of the injection site and swab with spirits. Treatment may be repeated at intervals of 24 hours.

## **WITHHOLDING PERIODS**

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residues Limits of Agricultural Compounds) Food Standards.**

Milk: Cattle - milk intended for sale for human consumption must be discarded during treatment and for the following times after the last treatment:

One dose – 48 hours (4 milkings);

two doses – 60 hours (5 milkings);

3 or more doses – 72 hours (6 milkings).

Sheep - milk – 35 days.

Meat: Animals producing meat and offal for human consumption must not be sold for slaughter either during treatment or within: sheep and cattle - 4 days; pigs - 5 days; horses - 63 days of the last treatment.

## **CONTRAINDICATIONS**

Not to be administered to animals known to be sensitive to penicillin. Occasional allergies to penicillin have been observed but these are very rare.

## **STORAGE**

Store in a cool place (2-8°C). Do not freeze. Short term (5 months) storage below 25°C is not expected to cause degradation of the product.

Registered pursuant to the ACVM Act 1997, No. A4256.

See [www.nzfsa.govt.nz/acvm/](http://www.nzfsa.govt.nz/acvm/) for registration conditions

Registered to:

Schering-Plough Animal Health Ltd

33 Whakatiki St, Upper Hutt, New Zealand.

Phone: 0800 800 543