

Keep out of reach of children
FOR ANIMAL TREATMENT ONLY

DUPLOCILLIN[®] LA

150,000 IU procaine penicillin and 150,000 IU benzathine penicillin per mL.

For S.C. or I.M. injection

100mL

Read the entire label before use

HANDLING PRECAUTIONS Warning contains penicillin which may cause an allergic skin reaction. Do not handle this product if you know you are sensitised, or if you have been advised not to work with such preparations. Use gloves, wash hands after use.

FIRST AID If on skin wash thoroughly with soap and water. If irritation or symptoms occur or persist, consult a doctor. Swelling of the face, eyes or lips or difficulty with breathing are more serious symptoms and require urgent medical attention. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately.

DISPOSAL Dispose of any unused contents and empty container in an approved landfill. Avoid release to the environment and avoid contamination of any water supply with product or empty container.

See safety data sheet for further information: www.intervet.co.nz

INDICATIONS

For the treatment of infections caused by bacteria sensitive to penicillin, where long-acting therapy is required, in horses, cattle, sheep, pigs, dogs and cats.

PROPERTIES

Procaine penicillin achieves a high initial penicillin blood level and the more slowly absorbed benzathine penicillin maintains a therapeutic blood level for 3-4 days.

CONTRAINDICATIONS

Not to be administered to animals known to be sensitive to penicillin. Occasional allergies to penicillin have been observed but these are rare.

DOSAGE

Large animals

12,000 IU penicillin per kg bodyweight. (1mL/25kg)

Companion animals (1mL/10kg)

30,000 IU penicillin per kg bodyweight.

e.g.

Horse	500kg	20mL
Cow	400kg	16mL
Sheep	50kg	2mL
Pig	50kg	2mL
Dog	10kg	1mL
Cat	5kg	0.5mL

ADMINISTRATION

By i.m. or s.c. injection. In food producing animals the injection is to be given into the anterior half of the neck.

WITHHOLDING PERIODS

Meat: Cattle, Sheep & Pigs 30 days – Animals producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 30 days of the last treatment. **Horses: 63 days**

Milk: Cattle 120 hours – Milk intended for sale for human consumption must be discarded during treatment and for not less than 10 milkings (120 hours) following the last treatment. **Sheep 35 days**
Not for use in bobby calves.

STORAGE

Store between 2-8°C. Do not freeze.

Prescription Animal Remedy (**P.A.R**) Class I.

For use only under the authority or prescription of a veterinarian.

Registered pursuant to the ACVM Act 1997, No. A4183.

See www.nzfsa.govt.nz/acvm/ for registration conditions.

Registered to:

Schering-Plough Animal Health Ltd

33 Whakatiki Street,

Upper Hutt, NZ.

Phone: 0800 800 543