

RESTRICTED VETERINARY MEDICINE

Keep out of reach of children
FOR ANIMAL TREATMENT ONLY

Nobivac[®] LEPTO I

LEPTOSPIROSIS VACCINE FOR DOGS

10 X 1 dose

DESCRIPTION

A liquid vaccine prepared from inactivated *Leptospira interrogans* Icterohaemorrhagiae, strain 820K.

INDICATIONS

Active immunisation against leptospirosis in dogs caused by *Leptospira interrogans* serovar Copenhageni.

DOSAGE AND ADMINISTRATION

Shake well before use. Allow to reach room temperature before using. Administer entire contents of the single dose vial by subcutaneous injection.

COMPATIBILITY

Nobivac Lepto 1 is compatible with and can be used to reconstitute the following Nobivac vaccines only:

- Nobivac DHPi
- Nobivac DHP
- Nobivac Parvo C
- Nobivac Puppy DP
- Nobivac Pi

VACCINATION PROGRAMME

Nobivac Lepto I should be used where there is a risk of dogs contracting leptospirosis. Can be used in pregnant bitches. The puppy programme can be completed from 10 weeks of age. The individual's age, health, breed and local environmental conditions should be considered before deciding on a vaccination programme.

Primary immunisation

Puppies should be at least 6 weeks old at time of first vaccination. Dogs and puppies should receive 2 doses, 2-4 weeks apart. An inter-dose interval of 4 weeks is preferable. The second dose should be given no earlier than 10 weeks of age.

Annual revaccination

A single dose is recommended.

	Nobivac Lepto I	Nobivac DHP	Nobivac KC
6-8 weeks	•	•	•
10 weeks	•	•	
1 year	•		•
2 year	•		•
3 year	•	•	•

PRECAUTIONS AND WARNING

- Only healthy dogs should be vaccinated.
- Sterile equipment should be used for administration.

ADVERSE REACTIONS

Following subcutaneous administration, a diffuse swelling up to 5mm in diameter may be observed at the site of injection. Occasionally this swelling may be hard and painful and last up to 3 days post injection. In rare cases, a transient rise in body temperature may occur.

Hypersensitivity reactions are rare, but as with all vaccines, may occasionally be encountered. In this event, administration of an antihistamine, corticosteroid or adrenaline by the most immediate route may be indicated.

STORAGE

- Stored at 2-8°C.
- Do not freeze.
- Protect from light.

ACVM Registration No. A7832.

See www.foodsafety.govt.nz for registration conditions.

Registered to:

Schering-Plough Animal Health Ltd

Phone: 0800 800 543

www.msd-animal-health.co.nz

Nobivac Parvo-C	ACVM No. A5525
Nobivac Puppy DP	ACVM No. A6018
Nobivac DHP	ACVM No. A5526
Nobivac Pi	ACVM No. A9101
Nobivac DHPPi	ACVM No. A6257

MSD Animal Health