

## **RESTRICTED VETERINARY MEDICINE**

Keep out of reach of children

FOR ANIMAL TREATMENT ONLY

# **Nobivac<sup>®</sup> TRICAT TRIO**

Live Attenuated Vaccine for Cats

**COMBINED FREEZE-DRIED FELINE HERPES VIRUS,  
CALICIVIRUS AND PANLEUCOPAENIA VIRUS VACCINE**

### **5 doses + diluent**

#### **DESCRIPTION**

A live freeze-dried vaccine, containing:

- Live attenuated feline calicivirus (strain F9)
- Live attenuated feline herpes virus type 1 (rhinotracheitis) (strain G2620A)
- Live attenuated feline panleucopaenia virus (strain MW-1)

It is presented as a freeze-dried plug; each pack contains 5 or 25 single dose vials. Pack also contains 5 or 25 corresponding 1mL vials of diluent for reconstitution prior to injection.

#### **INDICATION**

For active immunisation of cats from the age of 8-9 weeks onwards to:

- reduce clinical signs caused by an infection with feline calicivirus (FCV) and feline herpes virus type 1 (FHV)
- to reduce replication of feline calicivirus (FVC)
- and to prevent clinical signs, leucopenia and shedding caused by infection with feline panleucopaenia virus (FPLV)

The onset of immunity is 4 weeks for FCV & FHV, and 3 weeks for FPLV following first vaccination.

The duration of immunity is 1 year for the FCV and FHV components and 3 years for the FPLV component.

#### **DOSAGE AND ADMINISTRATION**

##### Initial programme

Two doses injected subcutaneously, at an interval of 3-4 weeks are required. The first vaccination is given from the age of 8-9 weeks and the second vaccination from the age of 12 weeks. If earlier vaccination is desirable or necessary, vaccination is appropriate from 6 weeks of age onwards, but should be followed by the normal vaccination scheme at 8-9 and 12 weeks of age.

##### Revaccination

A single dose injected subcutaneously.

Recommended revaccination following initial programme:

- FCV and FHV annual revaccination,
- FPLV revaccinate three-yearly.

#### **ADVICE ON CORRECT ADMINISTRATION**

Reconstitute the freeze-dried fraction of the vaccine with the accompanying diluent. Inject the diluent into the vial containing the freeze-dried pellet and shake gently until the pellet is completely reconstituted. Administer reconstituted vaccine by the subcutaneous route within 30 minutes of reconstitution. Use sterile injection equipment. Do not disinfect injection site.

#### **PRECAUTIONS**

Do not use during pregnancy or lactation, as the product has not been tested in pregnant or lactating queens.

## **ADVERSE REACTIONS**

Occasionally a slight painful swelling may be observed at the injection site for 1-2 days. A slight transient rise in body temperature (up to 40°C) may occur for 1-2 days. In some cases sneezing, coughing, nasal discharge and a slight dullness or reduced appetite may be observed for up to 2 days post vaccination. In very rare cases, the vaccine may cause hypersensitivity reactions (pruritus, dyspnoea, vomiting, diarrhoea and collapse).

Anaphylactic reactions are extremely rare, but as with all vaccines, may be encountered. In this event, administration of an antihistamine, corticosteroid or adrenaline by the most immediate route may be indicated.

## **SPECIAL WARNINGS**

Vaccinate only healthy animals.

An adequate clinical examination should be performed prior to vaccination.

Contact with potential sources of respiratory infection should be avoided until 7 days after the second vaccination.

Maternally derived antibodies, which may persist up to the age of 9-12 weeks, can have a negative influence on the efficacy of vaccination. In such cases, where a high level of maternally derived antibodies is expected, the vaccination schedule should be planned accordingly.

## **DISPOSAL**

Dispose of waste appropriately.

## **STORAGE**

Freeze-dried fraction: store at 2 - 8°C. Protect from light. Do not freeze

Diluent fraction: store between 2 - 25°C. Do not freeze.

## **FIRST AID**

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the doctor.

ACVM Registration No. A10164

See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions

Approval not required under the HSNO Act 1996.

Registered to:

**Schering-Plough Animal Health Ltd**

Phone 0800 800 543

[www.msd-animal-health.co.nz](http://www.msd-animal-health.co.nz)

