

Fact Sheet

Penalone®

Active Ingredient

Each syringe contains 1000mg procaine penicillin.

Indications

For the treatment of clinical mastitis in lactating cows.


How it works

Penicillin is the drug of choice to treat *Streptococcus uberis* (the most common mastitis pathogen isolated in New Zealand).

Penalone's unique formulation is designed to deliver a high dose of penicillin with good udder penetration, rapid absorption and long-lasting activity to achieve a first time cure. It is the obvious choice for early lactation mastitis, however excellent cure rates are also achieved throughout the entire lactation.

Efficacy

New Zealand trial data shows Penalone to be as effective as a penicillin – streptomycin combination intramammary with a proven record in the field (Streptopen HP)¹ throughout the entire lactation.



Dosage

Infuse the contents of 1 syringe into the infected quarter after each of 3 successive milkings (i.e. every 12 hours if milking twice daily).

Withholding period

Milk 9 milkings (108 hours if milking twice daily)

Meat 30 days

Pack size

20 Syringes

References

1. McDougall S, Weir A, Vermunt J, Bryan M, Arthur D. (2005) Clinical and bacteriological response to treatment of clinical mastitis with one of three intramammary antibiotics.

Prescription Animal Remedy (P.A.R.) Class I. For use only under the authority or prescription of a veterinarian. Registered pursuant to the ACVM Act 1997, No. A7795.
®Registered trademark. Schering-Plough Animal Health Limited, 33 Whakatiki Street, Upper Hutt, New Zealand. Phone 0800 800 543. PEN-309-2009