

Keep out of reach of children
FOR ANIMAL TREATMENT ONLY

Panacur[®] 100

Oral broad-spectrum drench for the management of internal parasites in cattle, sheep, goats, horses and deer.

Active ingredient: 100g/L fenbendazole

BROAD-SPECTRUM WHITE DRENCH

1 LITRE

Read entire label before use.

HANDLING PRECAUTIONS Warning may cause damage to the liver from repeated/prolonged oral exposure at high doses. Do not breath vapour.

FIRST AID If inhaled remove to fresh air. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor.

DISPOSAL Dispose of unused product through Agrecovery Chemicals. Triple rinse empty container, puncture and recycle through Agrecovery. Do NOT burn.

EMERGENCY RESPONSE For specialist advice in an emergency call 0800 CHEMCALL (0800 243 6225).

See Safety Data Sheet for further information: www.coopersonline.co.nz

Shake Well Before Use

PANACUR 100 is a free flowing white broad-spectrum oral anthelmintic for the treatment and control of mature, immature and egg stages of roundworms, including type II Ostertagia and lungworm. PANACUR 100 contains 100mg/mL of Fenbendazole.

ADMINISTRATION

By law the user must take due care, obtaining expert advice if necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

PANACUR 100 is given as an oral drench, and can be used through any standard drench gun.

RESISTANCE MANAGEMENT

PANACUR 100 contains fenbendazole, a member of the Benzimidazole family and is effective against internal parasites sensitive to this family of drenches.

- Resistance may develop to any chemical.
- Correct drenching techniques should be used.
- Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance.
- It is advisable that a resistance test be conducted regularly when using any parasite treatment.
- A representative sample of animals should be weighed before treatment.
- Dose the mob to the heaviest animal by liveweight in each group. Do not under dose.
- Check the accuracy of drench guns regularly.

WITHHOLDING PERIODS

It is an offence for the user of this product to cause residues exceeding the relevant MRL in the NZ (Maximum Residues Limits of Agricultural Compounds) Food Standards.

Meat: Sheep, goat, cattle and deer producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 10 days of the last treatment. Horses: 63 days.

Milk: Cattle: Milk intended for sale for human consumption must be discarded during treatment and for not less than 8 milkings (96 hours) following the last treatment.

Sheep & Goats: 35 days following the last treatment.

DOSE RATES

HORSES, CATTLE AND DEER	
0.75mL/10kg (7.5mg/kg)	
Liveweight	Dose Rate
50kg	4 mL
100kg	7.5mL
150kg	11.25mL
200kg	15 mL
250kg	19 mL
300kg	22.5mL
350kg	26.25 mL
400kg	30 mL
450kg	34 mL
500kg	37.5 mL
550kg	41.25 mL
600kg	45 mL
650kg	49 mL
Over 650kg	3.75mL/50kg liveweight

SHEEP AND GOATS	
0.5mL/10kg (5mg/kg)	
Liveweight	Dose Rate
40 kg	2 mL
50kg	2.5 mL
60kg	3 mL
70kg	3.5 mL
80kg	4 mL
Over 75kg	1 mL/20 kg liveweight

STORAGE

Store in a cool, dry place, avoiding temperatures below 0°C and above 30°C. **Do not allow to freeze.**

ACVM Registration No: A7154.

See www.nzfsa.govt.nz/acvm/ for registration conditions.

Approved pursuant to the HSNO Act 1996, HSR001924.

See www.ermanz.govt.nz for controls.

PANACUR and COOPERS are registered trademarks. © 2011 Schering-Plough Animal Health Limited.

Registered to:

Schering-Plough Animal Health Ltd

33 Whakatiki Street, Upper Hutt.

Phone: 0800 800 543

www.coopersonline.co.nz