

Keep out of reach of children
FOR ANIMAL TREATMENT ONLY

Penalone®

milking cow

SYRINGES

Schering-Plough Animal Health Limited

Each single dose syringe contains 1g procaine penicillin.

Read the entire label before use

HANDLING PRECAUTIONS Warning contains penicillin which may cause an allergic skin reaction. Do not handle this product if you know you are sensitised, or if you have been advised not to work with such preparations. Use gloves when handling this product to avoid exposure, wash hands after use.

FIRST AID In case of skin contact, carefully remove any contaminated clothing and wash skin thoroughly with soap and water. If irritation or symptoms occur or persist, consult a doctor. Swelling of the face, eyes or lips or difficulty with breathing are more serious symptoms and require urgent medical attention. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor.

DISPOSAL Dispose of contents according to label directions or in an approved landfill. **Container disposal** Dispose of empty syringe by burying in an approved landfill. Avoid release to the environment and avoid contamination of any water supply with product or empty container.

See Safety Data Sheet for further information. www.spah.co.nz.

USES

For the treatment of clinical mastitis caused by penicillin susceptible organisms in dairy cows during lactation.

DOSAGE

The contents of one syringe should be infused into the teat canal of each infected quarter immediately following each of three successive milkings, i.e. every 12 hours. Three syringes provide a full course of treatment.

ADMINISTRATION

Milk out the infected quarter. Clean teat thoroughly with cotton wool soaked in methylated spirit. Allow to dry. Infuse full contents of syringe into the teat canal of the infected quarter. Dip or spray teat with specifically formulated teat antiseptic.

1. After milking is complete, thoroughly clean and disinfect the end of the teat (e.g. with cotton wool soaked in methylated spirit).
2. Hold the barrel of the syringe firmly in one hand and gently twist and pull the protective cap to remove it. **DO NOT BEND THE NOZZLE.** Take care not to contaminate the nozzle.
3. Insert the nozzle into the teat canal and apply steady pressure on the syringe plunger until the full dose has been delivered. Holding the end of the teat in one hand, gently massage upwards with the other to aid dispersion of the antibiotic into the quarter.
4. Finally spray teat carefully with an approved teat spray.

WITHHOLDING PERIOD

MILK: Milk intended for sale for human consumption must be discarded during treatment and for not less than 9 milkings (108 hours) following the last treatment.

GUIDE TO HELP ACCURATE MILK DISCARD AFTER TREATMENT

Last treatment given	SUN AM	SUN PM	MON AM	MON PM	TUE AM	TUE PM	WED AM	WED PM	THU AM	THU PM	FRI AM	FRI PM	SAT AM	SAT PM
Milk may be added to the bulk tank	FRI AM	FRI PM	SAT AM	SAT PM	SUN AM	SUN PM	MON AM	MON PM	TUE AM	TUE PM	WED AM	WED PM	THU AM	THU PM

MEAT: Animals producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 30 days of the last treatment.

STORAGE

Store below 25°C. Do not refrigerate.

Prescription Animal Remedy (**P.A.R**) Class I.

For use only under the authority or prescription of a veterinarian.

Registered pursuant to the ACVM Act 1997, No. A7795.

See www.nzfsa.govt.nz/acvm/ for registration conditions.

Registered pursuant to the HNSO Act 1996, HSR002361.

See www.ermanz.govt.nz for controls.

PENALONE is a registered trademark of Schering-Plough Animal Health Limited
© 2006.

Registered to:

Schering-Plough Animal Health Limited

33 Whakatiki Street, Upper Hutt, New Zealand.

Phone: 0800 800 543. www.spah.co.nz