

## RESTRICTED VETERINARY MEDICINE

Keep out of reach of children.

FOR ANIMAL TREATMENT ONLY

# Piliguard<sup>®</sup> Pinkeye-1 Trivalent

PINKEYE VACCINE 100ML = 50 DOSES

*Moraxella Bovis* Bacterin

**Contains inactivated cultures of *Moraxella bovis* isolates in an oil emulsion adjuvant.  
Contains gentamicin as preservative.**

### INDICATION

For use in healthy cattle to aid in the control of pinkeye associated with infection by *Moraxella bovis* strains expressing pili similar to those expressed by isolates referred to as Strains Epp 63, Fla 64 and SAH 38.

### ADMINISTRATION

**Shake well before use.** The vaccine should be warmed to room temperature prior to injection. Inject 2mL intramuscularly into the anterior neck three to six weeks prior to onset of pinkeye season. Annual revaccination is recommended.

### CAUTION

Transient local reaction may occur at the injection site. Use may occasionally lead to development of granulomas which may persist for several weeks. Hypersensitivity reaction may occur and can cause temporary reduced milk production in lactating cattle. If anaphylaxis occurs administer epinephrine or antihistamine.

### STORAGE

Store at 2°C - 7°C. DO NOT FREEZE. Use entire contents when first opened.

### WARNING

Extreme caution should be used when injecting oil emulsion vaccines to avoid injecting yourself. Accidental injection can cause serious local reaction.

**Contact a doctor immediately if accidental injection occurs.**

ACVM Registration No. A8192.

See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.

Registered to and distributed in New Zealand by:

**Schering-Plough Animal Health Limited**

33 Whakatiki Street, Upper Hutt, New Zealand.

Phone 0800 800 543

[www.msd-animal-health.co.nz](http://www.msd-animal-health.co.nz)

