

Keep out of reach of children
FOR ANIMAL TREATMENT ONLY
YERSINIAVAX[®]

Protection against *Yersinia* losses in deer

Contains inactivated *Yersinia pseudotuberculosis* serotypes I, II & III.

100mL

Read entire label before use.

HANDLING PRECAUTION WARNING may cause an allergic skin reaction. Avoid skin contact.

FIRST AID If on skin, wash with plenty of soap and water. If skin irritation/rash occurs seek medical advice. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor.

ACCIDENTAL SELF-INJECTION Obtain medical attention - show this leaflet and/or SDS Accidental self injection may lead to an inflammatory response and medical advice should be sought on the management of deep injections, particularly those near a joint or associated with bruising. If possible the application of gentle squeezing pressure with absorbent material (e.g. facial tissues) at the injection site will swab up unabsorbed vaccine. Strong squeezing of the site should be avoided. The damaged area should be thoroughly cleansed and a topical antiseptic applied.

ADVICE TO DOCTOR Treat symptomatically. Some risk of hypersensitivity from injection. Contamination of the needle must be considered.

DISPOSAL Dispose of contents and empty vaxipak by burying in an approved landfill.

See Safety Data Sheet for further information. www.intervet.co.nz.

VACCINATION PROGRAMME

Which deer to vaccinate	1 st Dose	2 nd Dose
For use in young deer, at least 3 months of age.	At least 4-7 weeks before Yersiniosis is likely to occur.	3-6 weeks after the first dose. Allow 7-10 days for full protection to be attained.

DOSAGE

Inject a **2mL dose subcutaneously** in the anterior (front) half of the neck.

DIRECTIONS FOR USE

Shake well before use.

Equipment:

1. Ensure vaccinator delivers a **2mL dose**.
2. Attach end of draw-off tube to vaccinator. Pierce rubber bung of Vaxipak with draw-off needle.
3. Prime vaccinator as per vaccinator instructions.
4. Attach tie to Vaxipak and hang around neck.
5. Use sharp needles. Change needles approx every 20 deer.
6. Needles and vaccinator should be sterile prior to use.

Vaccination Technique:

1. **Vaccinate only healthy, clean, dry deer.**
2. Inject in the anterior (front) half of the neck.
3. Inject **subcutaneously i.e. between the skin and muscle.**
4. Raise skin to form a "tent" and insert needle into "tent opening" so that the needle is almost parallel with the neck.

DO NOT

Inject on the same side of the neck as TB tests.

SIDE EFFECTS

A site reaction or lesion may occur at the first injection site. It will normally disappear in 10-12 weeks. Deer should not be submitted for slaughter until the lesions have healed.

STORAGE

Store away from light at 2-8°C. Do not freeze. Unused vaccine must be discarded within 12 hours of opening.

Prescription Animal Remedy (**P.A.R**) Class I.

For use only under the authority or prescription of a veterinarian.

Registered pursuant to the ACVM Act 1997, No. A6151

See www.nzfsa.govt.nz/acvm/ for registration conditions.

Approved pursuant to the HSNO Act, 1996. Approval code HSR00015.

See www.ermanz.govt.nz for controls.

YERSINIAVAX and VAXIPAK are registered trademarks of Schering-Plough Animal Health Ltd.

Registered to:

Schering-Plough Animal Health Ltd

33 Whakatiki St,

Upper Hutt, New Zealand

Phone: 0800 800 543

www.intervet.co.nz